

Scattering in one-dimensional heterostructures described by the Dirac equation

N. M. R. Peres

Center of Physics and Department of Physics, University of Minho, P-4710-057, Braga, Portugal

Abstract. We consider electronic transport across one-dimensional heterostructures described by the Dirac equation. We discuss the cases where both the velocity and the mass are position dependent. We show how to generalize the Dirac Hamiltonian in order to obtain a Hermitian problem for spatial dependent velocity. We solve exactly the case where the position dependence of both velocity and mass is linear. In the case of velocity profiles, it is shown that there is no backscattering of Dirac electrons. In the case of the mass profile backscattering exists. In this case, it is shown that the linear mass profile induces less backscattering than the abrupt step-like profile. Our results are a first step to the study of similar problems in graphene.

PACS numbers: 72.10.-d, 73.21.Hb, 73.23.Ad

1. Introduction

Most of the accumulated knowledge about the physics of heterostructures assumes that the electrons in these materials are effectively described by the Schrödinger equation, with a position dependent mass. The common belief is that Dirac equation is of no use in condensed matter physics (spin-orbit coupling can be treated using the Pauli version of the Schrödinger equation). The discovery of graphene [1, 2] changed drastically this perspective. Condensed matter physicists are now facing a condensed matter system where the effective low-energy model for the quasiparticles is that of a ultrarelativistic, i. e. massless, Dirac equation, albeit with an effective Fermi velocity that is much lower than the velocity of light. In graphene the Fermi velocity is $v_F = c/300$, with c the velocity of light. In fact, since the isolation of graphene crystallites [1, 2, 3] that a renovated interest in the properties of the massless Dirac equation in 2+1 dimensions started to emerge [3, 4]. Although our interest in this paper is on the backscattering properties of electrons effectively described by the Dirac equation in 1+1 dimensions, as an effective low-energy theory of the electronic properties of a quasi-one-dimensional solid, we shall revise some of the properties of Dirac electrons in the context of graphene, given the accumulated amount of evidence for Dirac fermions in graphene.

How can the low-energy description of a solid be given by an effective Dirac-like equation? In one-dimensional physics, this takes place due to the linearization of the

spectrum close to the Fermi momentum, which introduces right- and left-movers with a linear energy-momentum relation. Another possibility, that we discuss in the bulk of the paper, is by having two atoms per unit cell with special type of hopping. In two-dimensions, this possibility has been realized by graphene. The electrons in graphene are confined to move within the π -orbitals of system, formed from the overlap of the $2p_z$ atomic orbitals of a single carbon atom. The transport and optical properties of graphene are determined essentially by the behavior of the π -electrons. The electronic density is such that graphene has one π -electron per carbon atom, and is therefore a half filled system. It turns out that the band structure of graphene around the *Fermi surface* is highly non-standard, since the relation between the momentum and the energy is linear, with a proportionality coefficient given by $v_F \simeq 1 \times 10^6$ m/s and termed Fermi velocity. In fact, the *Fermi surface* is reduced to two points (the so called Dirac points $\mathbf{K}$ and $\mathbf{K}'$) in the Brillouin zone, with a zero density of states at the Fermi-points. This fact gives graphene its non-conventional properties.

Starting from a $\mathbf{k} \cdot \mathbf{p}$ approach [5], adapted to degenerate bands off the Γ -point [6], we obtain an effective Hamiltonian in real space, valid around the $\mathbf{K}$ point, having the form

$$H = v_F \boldsymbol{\sigma} \cdot \mathbf{p} , \quad (1)$$

where $\boldsymbol{\sigma} = (\sigma_x, \sigma_y)$ is a vector using the Pauli matrices and $\mathbf{p} = -i\hbar(\partial_x, \partial_y)$. The Hamiltonian (1) is nothing but the Dirac Hamiltonian for massless particles in 2+1 dimensions, with an effective light-velocity v_F . Clearly, one expects that the physical properties of such a system will be different from those where the Schrödinger equation is valid. Note that this situation is an example of complex emergent behavior [12], since the original problem was that of independent electrons (and therefore Schrödinger like) in a periodic potential.

Those electrons in graphene lying close to the Fermi points ($\mathbf{K}$ and $\mathbf{K}'$), will have a different response to external potentials than those in other materials described by the Schrödinger equation. In fact, for Schrödinger electrons, if we consider that the potential varies at a given point in space from zero to a finite value V_0 (the so called step potential) there is always a finite fraction of impinging electrons that are reflected back. More over, if V_0 is greater than the energy of the impinging particles, there will be an exponential attenuated wave function in the region where the potential is finite.

For Dirac electrons the situation is different. It was shown by Klein [16] that Dirac particles can not be confined by an arbitrary large potential V_0 . In fact, they pass through strong repulsive potential without the exponential decay that characterizes Schrödinger particles. This is called Klein tunneling. If the particles are massless and moving in one-dimension the situation is even more dramatic, since there is no backscattered probability flux, no matter how large the potential is [13, 14].

In addition to external potentials, a particle propagating in solid state systems can face a situation where its effective mass changes in space. This possibility takes place in heterostructures, where the mass of the particle is written as $m(\mathbf{r})$, $\mathbf{r}$ being the position

of the particle. In this situation, the Schrödinger equation has to be modified, in order to comply with hermiticity and flux conservation.

If the particles are described by the Dirac equation, the information on the material properties is encoded both in the Fermi velocity [6] and in the mass. We can then imagine the possibility of producing a heterostructure where the Fermi velocity, $v_F(\mathbf{r})$, and the mass change with the position of the particle. As we show below, even in this case, there will be no backscattering if the particle is massless and the velocity is allowed to vary from point to point. In the context of graphene physics, a position-dependent Fermi velocity was already considered in the case of curved graphene [7]. Another possibility of producing both a Fermi velocity and a mass term that are position dependent is to subject the system to strain. It has already been shown using *ab-initio* calculations, in the context of graphene physics, that strain leads to gap formation [8, 9]. It is to be expected that for general strain, different parts of the material present local values of Fermi velocity and energy gap. Also in the case of graphene, it was shown that depositing the material on top of Boron Nitride leads to gap-formation [10].

In this paper we discuss several possibilities for the scattering of Dirac particles through a region where both the velocity and the mass are position dependent. Although our motivation is to study graphene strips [15] in a latter publication, we start by giving two exact solutions for one-dimensional systems. The generalization to a quasi-one-dimensional system, such as a narrow nano-wire, is easy to obtain and will be given in a follow up publication. In order to see the differences between the Schrödinger and the Dirac problems, we start by revisiting the case of one-dimensional heterostructures described by the Schrödinger equations and then move to the study of the Dirac case.

2. Warming up: the Schrödinger electrons in 1D

In order to compare how the scattering of Schrödinger electrons differs from that of Dirac particles, we give a brief account of the scattering of electrons by the interface of a heterostructure.

Let us consider the case of 1D Schrödinger electrons with position dependent effective mass $m(x)$, such that $m(x) = m^-$, for $x < 0$ and $m(x) = m^+$, for $x > 0$. The boundary conditions in this case are (see final paragraph of this section; for more general boundary conditions see Ref. [11])

$$\psi^-(0) = \psi^+(0), \quad (2)$$

$$\frac{1}{m^-} \frac{d}{dx} \psi^-(0) = \frac{1}{m^+} \frac{d}{dx} \psi^+(0). \quad (3)$$

We consider that the particles are moving from the left to the right. Therefore the wave function is

$$\psi^-(x) = e^{ikx} + r e^{-ikx}, \quad (4)$$

$$\psi^+(x) = t e^{ipx}, \quad (5)$$

where the values of k and p are fixed by energy conservation:

$$E = \frac{\hbar^2 k^2}{2m^-} = \frac{\hbar^2 p^2}{2m^+}. \quad (6)$$

The reflectance $|r|^2$ is determined from the boundary conditions as

$$|r|^2 = \frac{(m^+k - m^-p)^2}{(m^+k + m^-p)^2}, \quad (7)$$

which has the limiting value $|r|^2 \rightarrow 1$ for $m^+ \rightarrow \infty$. The transmittance coefficient $|t|^2$ is

$$|t|^2 = \frac{4(m^+k)^2}{(m^+k + m^-p)^2}, \quad (8)$$

We can check now that the boundary conditions used for the wave function are the correct ones, since they satisfy the flux current conservation. Using absolute values, we have indeed

$$j^{inc} = j^{refl} + j^{trans}, \quad (9)$$

with $j^{inc} = \hbar k/m^+$, $j^{refl} = |r|^2 \hbar k/m^+$, and $j^{trans} = |t|^2 \hbar p/m^-$.

The problem just described represents an heterostructure, where two materials are *glued* together (due to similar lattice constants), having different effective masses in the two materials. The used boundary conditions are obtained writing the Hamiltonian as

$$H = -\frac{\hbar^2}{2m} \frac{d}{dx} \frac{1}{m(x)} \frac{d}{dx}. \quad (10)$$

This choice defines an Hermitian problem $\ddagger$ since $\int dx (H\psi)^* \psi = \int dx \psi^* H\psi$. The construction of the (10) guaranties that the total probability flux is conserved. The boundary condition (3), follows immediately from the writting the Hamiltonian in the Sturm-Liouville form.

3. Dirac electrons in 1D

Let us now study the case of the Dirac Hamiltonian. We are interested in cases where both the velocity and the mass of the particles depend on their position in space. We study first the case of massless Dirac particles, like those present on graphene. Latter we add up the mass term. The Hamiltonian for a massless quasiparticle described by an effective Dirac equation, with an effective velocity of light v_F , is given by

$$H = v_F \sigma_x \frac{\hbar}{i} \frac{d}{dx}, \quad (11)$$

$\ddagger$ The Eq. (10) is a particular case of a Sturm-Liouville operator, which has the general form:

$$-\frac{d}{dx} \left(p(x) \frac{d}{dx} \right) + q(x)y = \lambda w(x)y,$$

where λ is to be determined from the boundary conditions, and $p(x)$, $q(x)$, and $w(x)$ are given functions. Sturm-Liouville problems are Hermitian.

where σ_x is the x Pauli matrix. Let us now consider that we make an heterostructure, made of two different materials, both of them described by an effective Dirac equation, such as (11). An example could be a strip of the material where part of it is subjected to strain and other part is strain free; this leads to different velocities in the two parts of the material. This situation calls for a model where v_F is position dependent: $v_F = v_F(x)$. Although this situation does not make sense in high-energy physics, it is quite conceivable in condensed matter physics, since the value of v_F is determined by the material under consideration – different materials can have different Fermi velocities.

The trivial replacement $v_F \rightarrow v_F(x)$ renders the problem non-Hermitian, as can be seen by applying the definition given above, when we discussed the Schrödinger case. There is however a way out. It is a simple matter to show that the operator

$$H = \sqrt{v_F(x)}\sigma_x \frac{\hbar}{i} \frac{d}{dx} \sqrt{v_F(x)}, \quad (12)$$

is Hermitian and reduces to Eq. (11) in the particular case $v_F(x) = v_F$. Note that the derivative will act on the product $\sqrt{v_F(x)}\psi(x)$. We stress that we are not studying the scattering of relativistic particles in condensed matter systems, but instead describing the scattering of particles represented by an effective low energy model that is formally equivalent to the Dirac equation (a situation that takes place in graphene). It would be interesting to derive Eq. (12) from a microscopic Hamiltonian. In Appendix A we give a tight-binding model whose effective low-energy theory is given by Eq. (12). The problem (12) has spinorial wave function of the form $\psi^\dagger = (\psi_1^*, \psi_2^*)$ and the probability flux is computed as $S_x = v_F(x)\psi^\dagger \sigma_x \psi$, as can be shown using the traditional derivation of computing the time change of the probability density [17].

The eigenproblem $H\psi = E\psi$, with H given by Eq. (12), corresponds to two coupled first-order differential equations of the form

$$\sqrt{v(x)} \frac{\hbar}{i} \frac{d}{dx} [\sqrt{v(x)}\psi_2(x)] = E\psi_1(x), \quad (13)$$

$$\sqrt{v(x)} \frac{\hbar}{i} \frac{d}{dx} [\sqrt{v(x)}\psi_1(x)] = E\psi_2(x). \quad (14)$$

It is straightforward to show that the two first-order differential equations given above can be put in Sturm-Liouville form:

$$-\frac{d}{dx} [v_F(x) \frac{d}{dx} y_1(x)] = \frac{E^2}{v_F(x)\hbar^2} y_1(x), \quad (15)$$

with $y_1(x) = \sqrt{v_F(x)}\psi_1(x)$, $p(x) = v_F(x)$, $\lambda = E^2/\hbar^2$, and $w(x) = 1/v_F(x)$, and $\psi_2(x)$ obtained from

$$\psi_2(x) = \frac{\hbar}{iE} \sqrt{v_F(x)} \frac{d}{dx} [\sqrt{v_F(x)}\psi_1(x)]. \quad (16)$$

If the velocity profile changes continuously, one expects that the continuity of the wave function should be valid. If the velocity profile changes abruptly at a given point in space, say at $x = 0$, from $v(x) = v_-$, for $x < 0$, to $v(x) = v_+$, for $x > 0$, one has to use

Eqs. (13) and (14) to derive the boundary conditions. Defining $y_2(x) = \sqrt{v_F(x)}\psi_2(x)$ Eqs. (13) and (14) can be written as

$$\frac{\hbar}{i} \frac{d[y_2(x)]}{dx} = E \frac{y_1(x)}{v(x)}, \quad (17)$$

$$\frac{\hbar}{i} \frac{d[y_1(x)]}{dx} = E \frac{y_2(x)}{v(x)}. \quad (18)$$

Integrating Eqs. (17) and (18) using a symmetric infinitesimal interval around $x = 0$ one obtains the condition

$$y_i(0^-) = y_i(0^+), \quad (19)$$

with $i = 1, 2$. Note that condition (19) implies the discontinuity of the wave function. Let us now move to the solution of several particular cases.

3.1. The step-like velocity profile

Let us consider the case

$$v_F(x) = v_F^- \theta(-x) + v_F^+ \theta(x). \quad (20)$$

The solution of the Dirac equation reads

$$\psi_-(x) = \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{iqx} + r \begin{pmatrix} 1 \\ -1 \end{pmatrix} e^{-iqx}, \quad (21)$$

and

$$\psi_+(x) = t \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{ipx}, \quad (22)$$

with $E = v_F^- q \hbar$ and $E = v_F^+ p \hbar$. The above wave functions were obtained by solving the Dirac equation for $x \gtrless 0$, where the velocity is constant. Using these solutions and the boundary condition (19) one obtains

$$\begin{cases} \sqrt{v_-}(1+r) = \sqrt{v_+}t, \\ \sqrt{v_-}(1-r) = \sqrt{v_+}t, \end{cases} \quad (23)$$

which is satisfied only for $r = 0$. Had we consider the general case of both a velocity profile (20) and a potential profile of the form

$$V(x) = V_0 \theta(x), \quad (24)$$

under the condition $E > V_0$, the boundary conditions would still give $r = 0$. This result is called Klein tunnelling [16]. If we consider the same velocity and potential profiles, but now take the energy $E < V_0$, the boundary conditions still gives the result (23), but the wave function of the propagating mode for $x > 0$ is now different and given by

$$\psi_+(x) = t \begin{pmatrix} 1 \\ 1 \end{pmatrix} e^{-ipx}. \quad (25)$$

In this case the result will also be $r = 0$. The conclusion is that it is not possible to backscatterer massless Dirac electrons with a step-like velocity and potential profiles in 1D.

3.2. The linear velocity profile with massless particles

We have seen that an abrupt change of the velocity at the interface produces no reflected particles. Let us now study the case of a smooth change in the velocity from v_F^- to v_F^+ . (The result can be guessed from the outset!) To that end, we choose the profile

$$v_F(x) = v_F^- \theta(-\delta - x) + (\bar{v} + x\Delta) \theta(\delta - |x|) + v_F^+ \theta(x - \delta), \quad (26)$$

where we have defined $\bar{v}$ and Δ as

$$\bar{v} = \frac{v_F^- + v_F^+}{2}, \quad (27)$$

$$\Delta = \frac{v_F^+ - v_F^-}{2\delta}. \quad (28)$$

For the cases $x < -\delta$ (region *I*) or $x > \delta$ (region *III*) the Fermi velocity is constant and the solution of the Dirac equation is elementary, as we have seen before. The interesting case is therefore the region $|x| < \delta$ (region *II*). In this case we have to use the Dirac equation in the form (12). Explicitly we have to solve the problem $H\psi = E\psi$ with H given by

$$H = \sqrt{\bar{v} + x\Delta} \sigma_x \frac{\hbar}{i} \frac{d}{dx} \sqrt{\bar{v} + x\Delta}. \quad (29)$$

Writting the differential equations satisfied by the spinors, we obtain for the ψ_1 spinor (considering the substitution $y = \sqrt{\bar{v} + x\Delta}\psi_1$) the equation

$$-v_F^2(x) \frac{d^2 y}{dx^2} - \Delta v_F(x) \frac{dy}{dx} = \epsilon^2 y, \quad (30)$$

with $\epsilon^2 = E^2/\hbar^2$. Making the replacement

$$z = \frac{\bar{v} + x\Delta}{\bar{v}} \equiv \theta(x), \quad (31)$$

we obtain the result

$$z^2 \frac{d^2 y}{dz^2} + z \frac{dy}{dz} + \nu^2 y = 0, \quad (32)$$

with $\nu^2 = \epsilon^2/\Delta^2$. Making the additional replacement $\omega = \ln z$, Eq. (32) is reduced to that of the harmonic oscillator §

$$\frac{d^2 y}{d\omega^2} + \nu^2 y = 0. \quad (33)$$

The general solution of Eq. (33) is elementary and from it one obtains $y(x)$ given by

$$y(x) = A \sin[\nu L \theta(x)] + B \cos[\nu L \theta(x)], \quad (34)$$

with

$$L\theta(x) = \ln \frac{\bar{v} + x\Delta}{\bar{v}}. \quad (35)$$

§ One should note that the solution of Eq. (32) can also be obtained by noticing that $y = z^m$ is a solution if $m^2 = -\nu^2$, leading to $y(z) = Az^{i\nu} + Bz^{-i\nu}$.

The component ψ_1 of the spinor is obtained from $\psi_1(x) = y(x)/\sqrt{v_F(x)}$, and ψ_2 is obtained using Eq. (16) and reads

$$\psi_2(x) = \frac{\hbar}{iE} \sqrt{v_F(x)} \frac{dy(x)}{dx}. \quad (36)$$

The boundary conditions are $\psi^I(-\delta) = \psi^{II}(-\delta)$ and $\psi^{II}(\delta) = \psi^{III}(\delta)$, and can be written as ||

$$\begin{pmatrix} -e^{ik\delta} & S_-/\sqrt{v_F^-} & C_-/\sqrt{v_F^-} & 0 \\ e^{ik\delta} & C_- \lambda^- & -S_- \lambda^+ & 0 \\ 0 & S_+/\sqrt{v_F^+} & C_+/\sqrt{v_F^+} & -e^{iq\delta} \\ 0 & C_+ \lambda^- & -S_+ \lambda^+ & -e^{iq\delta} \end{pmatrix} \begin{pmatrix} r \\ A \\ B \\ t \end{pmatrix} = \begin{pmatrix} e^{-ik\delta} \\ e^{-ik\delta} \\ 0 \\ 0 \end{pmatrix} \quad (37)$$

with

$$S_{\pm} = \sin[\nu L \theta(\pm\delta)], \quad (38)$$

$$C_{\pm} = \sin[\nu L \theta(\pm\delta)], \quad (39)$$

$$\lambda_{\pm} = \frac{\hbar \sqrt{v_F^{\pm}} \Delta}{iE} \frac{\nu}{\bar{v} \theta(\pm\delta)}. \quad (40)$$

The fraction of reflected flux is given by $|r|^2$ and the transmitted flux is $1 - |r|^2 = v_F^+ |t|^2 / v_F^-$. The explicit evaluation (which is some what lengthy) of the coefficients gives

$$|r|^2 = 0, \quad (41)$$

$$|t|^2 = \frac{v_F^-}{v_F^+}, \quad (42)$$

and therefore the electrons are totally transmitted across the heterojunction. Of course that this result could have been anticipated from the conclusions of Sec. 3.1, since it is always possible to represent a well-behaved function by a sum of infinitesimal rectangles. (It is however elegant to have an exact solution to a given problem.)

3.3. The linear velocity profile with massive particles

We can now use what we have just learned to deal with a more general situation where the electronic spectrum changes due to a change of the material. The simplest case is that where for $x = \delta$ the mass of the quasi-particles jumps from zero to a finite value. The connection between the two materials is represented by the linear profile of the velocity discussed in Sec. 3.2. The mass profile is

$$m(x) = m \theta(x - \delta), \quad (43)$$

that is, the change in velocity takes place outside the region of finite mass. Since the particle has a finite mass the spectrum changes to $E = \pm \sqrt{v_+^2 q^2 + m^2 v_+^4}$, and the wave function in this region changes to

$$\psi^{III}(x) = t \begin{pmatrix} \sqrt{E^2 - m^2 v_+^4} \\ E - m v_+^2 \end{pmatrix} e^{-iqx}. \quad (44)$$

|| Note that the dimensions of A and B are $\sqrt{L/T}$, and $\lambda^{\pm}$ has dimensions of $\sqrt{T/L}$.

The overall modification is the replacement of e^{iq} by $\sqrt{E^2 - m^2 v_+^4} e^{iq}$ in the third row, and of e^{iq} by $(E - mv_+^2) e^{iq}$ in the fourth row. Working out the calculation of $|r|^2$ we obtain

$$|r|^2 = -[1 + 2\epsilon(-\epsilon + \sqrt{\epsilon^2 - 1})], \quad (45)$$

with $\epsilon = E/(mv_+^2)$. As expected, the result does not depend on δ and on $v_F^\pm$, due to the reasons discussed in Sec. 3.2. A plot of Eq. (45) is given in Fig. 1. The situation is same if the mass profile had been chosen as

$$m(x) = m\theta(x + \delta), \quad (46)$$

this is, the linear profile




Figure 1. Left: plot of Eq. (45) as function of ϵ . Right: plot of Eq. (45) as function of ϵ

3.4. The linear mass profile

We have seen above that the linear velocity profile does not contribute to the backscattering of Dirac-fermions, even when it is combined with a region where fermions are massive. In fact, any velocity profile produces no backscattering in one-dimension. All the backscattering comes from the change in the dispersion, due to the presence of the mass term. This fact motivates the question of what is form of the wave function and of the coefficient $|r|^2$ if the mass does not change abruptly but in a smooth way? A choice for the change of the mass profile, leading to an exact solution, is that described by:

$$m(x) = m\theta(x - \delta) + m\frac{x}{\delta}\theta(x)\theta(\delta - x). \quad (47)$$

We approach the solution of the scattering problem by solving first the Dirac equation subjected a mass profile $m(x) = m\frac{x}{\delta}$. The method of solution is inspired in that used for the 3+1 Dirac equation [18, 19]. In this case the Fermi velocity is constant, and therefore we have to solve Eq. (11) with the additional term $\sigma_z m v_F^2 \frac{x}{\delta} \psi$:

$$v_F \sigma_x \frac{\hbar}{i} \frac{d\psi}{dx} + \sigma_z m v_F^2 \frac{x}{\delta} \psi = E \psi. \quad (48)$$

The solution of this problem proceeds in several steps. The first is to operate on the left of Eq. (48) with the operator

$$\frac{d}{dx}\sigma_y\sigma_z. \quad (49)$$

After some algebra we obtain the following result

$$\hbar^2 v_F^2 \frac{d^2 \psi}{dx^2} = -[(E^2 - m^2 v_F^4 x^2 / \delta^2) + \sigma_y \hbar m v_F^3 / \delta] \psi. \quad (50)$$

The two components of the spinor are still coupled in Eq. (50). In order to decouple them we use the unitary transformation

$$\phi = U^\dagger \psi, \quad (51)$$

$$U = \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}. \quad (52)$$

The above transformation changes σ_y to $\tilde{\sigma}_y = U^\dagger \sigma_y U$, which, still mixing the spinors in the Eq. (53) below, does it with the same sign (this is a crucial step). After applying the unitary transformation we obtain

$$\hbar^2 v_F^2 \frac{d^2 \phi}{dx^2} = -[(E^2 - m^2 v_F^4 x^2 / \delta^2) + \tilde{\sigma}_y \hbar m v_F^3 / \delta] \phi. \quad (53)$$

We now introduce two new functions defined by $F_\pm = \phi_1 \pm \phi_2$, where ϕ_i (with $i = 1, 2$) are the components of the spinor ϕ . In terms of the functions $F_\pm$ the eigenproblem takes the form:

$$-\hbar^2 v_F^2 \frac{d^2 F_\pm}{dx^2} + [m^2 v_F^4 x^2 / \delta^2 - \epsilon_\pm^2] F_\pm = 0, \quad (54)$$

where $\epsilon_\pm^2 = E^2 \pm \hbar m v_F^3 / \delta$. The general solution of Eq. (54) is given in terms of parabolic cylinder functions $D_\nu(x)$ (see Appendix B)

$$F_\pm(x) = A_\pm D_{\nu_1^\pm}(\sqrt{2}(b/a)^{1/4}x) + B_\pm D_{\nu_2^\pm}(i\sqrt{2}(b/a)^{1/4}x), \quad (55)$$

with

$$\nu_1^\pm = -\frac{1}{2} + \frac{\epsilon_\pm^2}{2\sqrt{ab}} = \nu - (1 \mp 1)/2, \quad (56)$$

$$\nu_2^\pm = -\frac{1}{2} - \frac{\epsilon_\pm^2}{2\sqrt{ab}} = -\nu_1^\pm - 1 \quad (57)$$

$$\nu = \frac{E^2 \delta}{2\hbar v_F^3 m}, \quad (58)$$

$$a = \hbar^2 v_F^2, \quad (59)$$

$$b = v_F^4 m^2 / \delta^2. \quad (60)$$

If we were looking for the solution of the problem valid to all values of x , the functions $D_\nu(x)$ with imaginary argument, since are not real and are not normalizable in the infinite volume, had to be excluded. Therefore, in the limit $x \rightarrow \pm\infty$ the normalizable

solutions are those where $F_{\pm}(x)$ represents the one-dimensional harmonic oscillator wave functions (see Appendix C). Therefore the solution in the interval $x \in [-\infty, \infty]$ is

$$F_{\pm}(x) = 2A_{\pm}D_{\nu_{\pm}^{\pm}}(\sqrt{2}(b/a)^{1/4}x). \quad (61)$$

This choice guaranties that the wave function is well behaved at $x \rightarrow \pm\infty$ when $\nu_1^{\pm}$ is equal to a positive integer (see Appendix C). Therefore, in the infinite volume, the solution of Eq. (50) takes the form

$$\psi(x) = \begin{pmatrix} A_+D_{\nu_1^+} + A_-D_{\nu_1^-} \\ iA_+D_{\nu_1^+} - iA_-D_{\nu_1^-} \end{pmatrix}, \quad (62)$$

where the argument of the parabolic cylinder functions has been omitted for simplicity of writing. We still have to check whether Eq. (62) is solution of Eq. (48). Introducing the solution (62) in Eq. (48) we obtain that the wave function is a solution if

$$A_- = \frac{\Lambda\nu}{E}A_+, \quad (63)$$

$$\Lambda = v_F\hbar\sqrt{\frac{2v_F^2m}{v_F\hbar\delta}} = v_F\hbar/\beta. \quad (64)$$

What we have discuss so far assumes that Eq. (48) holds for every x . Our interest, however, is on the scattering problem of electrons when Eq. (48) holds for $x \in]0, \delta[$. In this case, the general solution (55) with both real and complex wave functions must be used. The strategy is the same used before. We known at the outset that the solution of Eq. (48) is obtained from the solution (62) by fixing the value of a constant, as in Eq. (63). The final solution is

$$\begin{aligned} \psi(x) = & A \begin{pmatrix} D_{\nu}(x/\beta) + \nu\Lambda/ED_{\nu-1}(x/\beta) \\ iD_{\nu}(x/\beta) - i\nu\Lambda/ED_{\nu-1}(x/\beta) \end{pmatrix} \\ & + B \begin{pmatrix} D_{-\nu}(ix/\beta) + i\nu\Lambda/ED_{-\nu-1}(ix/\beta) \\ -iD_{-\nu}(ix/\beta) - \nu\Lambda/ED_{-\nu-1}(ix/\beta) \end{pmatrix} \end{aligned} \quad (65)$$

which is valid in the region $x \in]0, \delta[$. As in Sec. 3.2, we wave the incoming wave function given by Eq. (21), for $x < 0$, and the transmitted one given by Eq. (44), for $x > \delta$. The reflected and transmitted amplitudes r and t , respectively, are obtained from imposing the continuity of the wave function at $x = 0$ and $x = \delta$. In Figure 2 we give some numerical examples of our calculation. It is clear from Fig. 2 that the smoothness of the change of the mass profile leads to a larger transmittance, for a given energy, than when the mass-profile changes abruptly. Also, when the change in the value of the mass takes place over a relatively large regions, the transmittance shows the presence of resonances, as can be seen in panel (c) of Fig. 2.

4. Discussion and conclusions

We have studied several scattering problems using a modified version of the Dirac Hamiltonian which incorporates the possibility of a spatial dependent velocity and mass terms. Two exact solutions were given. We showed that in the case of a velocity profile it is necessary the modification of the original Dirac Hamiltonian, in order to have a Hermitian problem. We have shown that Klein tunneling is not suppressed by a change on the velocity profile, with a transmittance equal to unity always. This was understood by studying the case of an abrupt change in the velocity profile and also by solving exactly the case where the velocity changes linearly across a given region.

We have also studied the case where the mass term depends on position. For this situation, we solved the cases of an abrupt change of the mass value and of a linear change of the mass profile. In both cases we see the presence of backscattering, with values of the transmittance smaller than unity. The smoother mass profile induces less amount of backscattering.

It is interesting to consider next the case of Dirac electrons in a strip of finite width W , a situation relevant for graphene strips, and see if in this case a position dependent velocity profile does produce backscattering. This will be considered in a follow up publication.

Acknowledgments

This work was supported by FCT under the grant PTDC/FIS/64404/2006. J. M. B. Lopes dos Santos and Vitor Pereira are acknowledged for fruitful discussions. J. L. Martins is acknowledged for suggesting the form of the Hermitian Dirac Hamiltonian.




Figure 2. Transmittance of Dirac electrons across a region where the mass profile changes according to Eq. (47). The dotted line represents the transmittance in the case of an abrupt change of the mass profile, as given by Eq. (45). The energy gap is represented by the value of mv_F^2 . In panel (a) we give the results for a value of δ of 1 nm. In panels (b) and (c) we plot the same but for $\delta = 10$ nm. In panel (c) we give a zoom in of the solid curve of panel (b); it is clear the existence of resonances in the transmittance. The Fermi velocity is 10^6 m/s, leading to $v_F \hbar = 0.66$ eV·nm.

Appendix A. A derivation of the Dirac equation with a position dependent velocity

Consider a one-dimensional tight-binding Hamiltonian with two atoms per unit cell such that the hopping parameter within a unit cell n is $-t_n$ and among nearest unit cells in t_n . The Hamiltonian in second quantization reads

$$H = \sum_n [-t_n(a_n^\dagger b_n + b_n^\dagger a_n) + t_n(a_n^\dagger b_{n-1} + b_{n-1}^\dagger a_n)]. \quad (\text{A.1})$$

The in the case $t_n = t$ (the hopping is independent of the unit cell) the spectrum of the electrons is

$$E_{\alpha=\pm} = \alpha 2t |\sin(kc/2)|, \quad (\text{A.2})$$

with c is the length of the unit cell vector. Close to $k = 0$ (zero energy) the spectrum is linear in momentum and a massless Dirac spectrum is generated with a Fermi velocity given by v_F . Let us now assume the general case of a t_n dependent on the unit cell position and obtain from the Hamiltonian (A.1) the effective field-theoretical model that describes the system at low energy ($E \simeq 0$). Since the momentum close to which the Dirac spectrum develops is $k = 0$, we can write immediately the effective field theoretical model as

$$H = -\frac{1}{c} \int dy t(y) \left[a^\dagger(y) c \frac{db(y)}{dy} + c \frac{db^\dagger(y)}{dy} a(y) \right], \quad (\text{A.3})$$

where we have used the expansion $b^\dagger(y-c) \simeq b^\dagger(y) + cd[b^\dagger(y)]/dy$. Integrating by parts the second term of Eq. (A.3) and using the Pauli matrices to help condensing the results we obtain

$$H = \frac{1}{c} \int dy \left[\Psi^\dagger \sqrt{v_y(y)} \sigma_y p_y \sqrt{v_y(y)} \Psi + \Psi^\dagger \frac{c\sigma_x}{2} \frac{dt(y)}{dy} \Psi \right] \quad (\text{A.4})$$

with $v(y) = ct(y)/\hbar$ and $\Psi^\dagger = [a^\dagger(y)b^\dagger(y)]$. The first term in Eq. (A.4) is our proposed Hamiltonian (12).

Appendix B. Weber's differential equation

We give here some basic information on the Weber's differential equation, aiming to give the text a selfcontain nature and to fix notation and definitions. Weber's differential equation is defined as

$$y''(z) + (\nu + 1/2 - z^2/4)y(z) = 0, \quad (\text{B.1})$$

and its two independent solutions are the parabolic cylinder functions $y(z) = D_\nu(z)$ and $y(z) = D_{-\nu-1}(iz)$. Equation (54) is of the general form

$$y''(x) + (-ax^2 + c)y(x) = 0. \quad (\text{B.2})$$

Making the transformation $x = \beta z$, with β given by $\beta = (4a)^{-1/4}$ and $\nu = -1/2 + c\beta^2$, we reduce Eq. (B.2) to Weber's equation (B.1). Using for a and c the particular values of our problem we obtain

$$\beta = \sqrt{\frac{\hbar v_F \delta}{2v_F^2 m}}, \quad (\text{B.3})$$

and

$$\nu = -\frac{1}{2} + \frac{\epsilon_{\pm}^2 \delta}{2\hbar v_F^3 m}. \quad (\text{B.4})$$

The derivative of the parabolic cylinder functions obeys

$$D'_\nu(z) + zD_\nu(z)/2 - \nu D_{\nu-1}(z) = 0. \quad (\text{B.5})$$

$$D_{\nu+1}(z) - zD_\nu(z) + \nu D_{\nu-1}(z) = 0. \quad (\text{B.6})$$

Using the results of Ref. [[20]], the solution of $D_\nu(z)$ can be written in terms of the Kummer confluent hypergeometric function, $U(a, b, x)$, as

$$D_\nu(z) = 2^{\nu/2} e^{-z^2/4} U(-\nu/2, 1/2, z^2/2). \quad (\text{B.7})$$

The Kummer function $U(a, 1/2, z)$ is computed using the the Kummer confluent hypergeometric function, $M(a, b, x)$, as

$$U(a, 1/2, z) = \sqrt{\pi} \frac{M(a, 1/2, z)}{\Gamma(a + 1/2)} - 2\sqrt{z\pi} \frac{M(a + 1/2, 3/2, z)}{\Gamma(a)}. \quad (\text{B.8})$$

Appendix C. Eigenvalues of the scalar potential $V(x) = \sigma_z m v_F^2 x / \delta$

The problem we introduced in Sec. 3.4 was that of a particle that moves in a heterostructure with a mass dependent position. We can, however, think of this problem as that of a Schrödinger particle moving in the scalar potential $V(x) = m^2 v_F^4 (x/\delta)^2$. Since Eq. (54) is that of an one-dimensional harmonic oscillator, the normalizable solutions have the well known form

$$z_n(u) = \frac{\pi^{-1/4}}{\sqrt{2^n n!}} e^{-u^2/2} H_n(u), \quad (\text{C.1})$$

where

$$u = x \left(\frac{V_F^2 m}{\delta V_F \hbar} \right)^{1/2} \quad (\text{C.2})$$

upon the identification

$$\frac{1}{m_0} \leftrightarrow 2v_F^2, \quad (\text{C.3})$$

$$\omega_0^2 \leftrightarrow 4v_F^6 m^2 / \delta^2, \quad (\text{C.4})$$

with m_0 and ω_0 the mass and the frequency of the oscillator, respectively. The spectrum is obtained from $\epsilon_{\pm}^2 = 2\hbar v_F^3 m(n + 1/2)/\delta$, with $n = 0, 1, 2, \dots$, leading to

$$E_+^2 = 2\hbar v_F^3 m n \delta^{-1}, \quad (\text{C.5})$$

$$E_-^2 = 2\hbar v_F^3 m(n + 1) \delta^{-1}. \quad (\text{C.6})$$

Since the energy has to be same for both F_+ and F_- , we choose the solutions $\P F_+^n = z_n$ and $F_-^n = z_{n-1}$, with $z_{-1} = 0$. Finally, from the definition $F_{\pm} = \phi_1 \pm \phi_2$, we obtain (properly normalized)

$$\phi_1(x) = [z_n(x) + z_{n-1}(x)]/\sqrt{2}, \quad (\text{C.7})$$

$$\phi_2(x) = [z_n(x) - z_{n-1}(x)]/\sqrt{2}. \quad (\text{C.8})$$

The above equations are the solution of Eq. (50).

References

- [1] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, *Science* **306**, 666 (2004).
- [2] K. S. Novoselov, D. Jiang, T. Booth, V.V. Khotkevich, S. M. Morozov, A. K. Geim, *PNAS* **102**, 10451 (2005).
- [3] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *Review of Modern Physics* (in print); arXiv:0709.1163.
- [4] Andreas Barth and Werner Marx, arXiv:0808.3320.
- [5] John H. Davies, *The physics of low-dimensional semiconductors* (New York: Cambridge University Press, 1998).
- [6] D. P. DiVincenzo and E. J. Mele, *Phys. Rev. B* **29**, 1685 (1984).
- [7] Fernando de Juan, Alberto Cortijo, and María A. H. Vozmediano, *Phys. Rev. B* **76**, 165409 (2007).
- [8] G. Gui, J. Li, and J. Zhong, *Phys. Rev. B* **78**, 075435 (2008).
- [9] Zhen Hua Ni, Ting Yu, Yun Hao Lu, Ying Ying Wang, Yuan Ping Feng, and Ze Xiang Shen, arXiv:0810.3476.
- [10] Gianluca Giovannetti, Petr A. Khomyakov, Geert Brocks, Paul J. Kelly, and Jeroen van den Brink, *Phys. Rev. B* **76**, 73103 (2007).
- [11] W. A. Harrison and Alexander Kozlov, *Matching conditions in effective mass theory*. In Ping Jiang and Hou-Zhi Zheng, editors, *Proceedings of the 21st International Conference on the Physics of Semiconductors*, (Singapore: World Scientific, 1992) p. 341.
- [12] P. W. Anderson, *Science* **177**, 393 (1972).
- [13] M. I. Katsnelson, K. S. Novoselov, and A. K. Geim, *Nature Physics* **2**, 620 (2006).
- [14] C. W. Beenakker, arXiv:0710.3848.
- [15] J. Viana Gomes and N. M. R. Peres, *J. Phys.: Cond. Matt.* **20**, 325221 (2008)
- [16] A. Calogeracos and N. Dombey, *Contemp. Phys.* **40**, 313 (1999).
- [17] Leonard I. Schiff, *Quantum Mechanics*, 3^{ed} (Singapore: McGraw-Hill, 1968).
- [18] G. Calucci, *Lett. Nuovo Cimento* **26**, 449 (1979).
- [19] Long Chao-Yun and Qin Shui-Jie, *Chinese Physics* **16**, 897 (2007).
- [20] Milton Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, (New York: Dover, 1965).

$\P$ It is interesting to note that the system supports a zero energy mode: $F_+^{n=0} = z_0$ and $F_-^{-1} = 0$.